

CS105L: Discrete Structures

I semester, 2005-06

Homework # 6

Due before class on **Tuesday, October 4, 2005**

Instructor: Amitabha Bagchi

September 26, 2005

1. A *vertex colouring* of a graph is an assignment of colours (or just natural numbers) to the vertices of a graph. A *colour class* in a coloured graph is a maximal set of vertices which have been assigned the same colour. A graph is said to be (k, d) -colourable if the vertices can be coloured with k colours in such a way that the vertices in each colour class induce a graph of maximum degree d . Prove the following theorem due to Lovász:

Theorem 6.1 (Lovász, 1966) *For any k , any graph of maximum degree Δ with $|E|$ edges can be $(k, \lfloor \Delta/k \rfloor)$ -coloured in time $O(\Delta|E|)$.*

2. Let us consider a multigraph G . To avoid confusion let us invent the term *edge-holder* which means a vertex pair and let us use the term *edge* to mean an actual edge between two vertices. In a multigraph the set of edge-holders is a set, and the set of edges is a multiset chosen from the set of edge-holders. In fact if there is no edge corresponding to a given edge-holder, then that edge-holder is of no interest to us. With this in mind let us try to prove a generalization of Menger's theorem (the edge version.)

Given a pair of vertices (s, t) in a graph G , let us consider a maximal set of edge disjoint paths P between s and t (note, the paths of P are not edge-holder disjoint.) We call such a set of paths *k-balanced* if every edge holder occurs in less than $|P|/k$ paths of P . If there is an edge holder which occurs exactly $\lfloor |P|/k \rfloor$ times in P then it is known as a *critical edge*.

- (a) Prove, using Menger's theorem, that if a set of edge-disjoint paths P between s and t is k -balanced then it is possible to find a set of k edge-holder disjoint paths between s and t using the edges in P .
- (b) Prove that it is possible to find $\lfloor |P|/k \rfloor$ sets of k -edge holder disjoint paths between s and t using the edges in P . Note that each set of k paths has to be edge holder disjoint within itself but might share edge holders with other such sets.

To prove the second part you will need the following lemma:

Lemma 6.1 *Given a k -balanced path system with critical edges between two vertices s and t , there is a set of k -edge holder disjoint paths from s to t which pass through every critical edge.*