

## TUTORIAL SHEET 10

1. Suppose you have algorithm  $A$  which given a graph  $G$  and a number  $k$ , outputs YES iff  $G$  has a vertex cover of size at most  $k$ . Assuming that  $A$  runs in polynomial time, show that you can find a vertex cover of minimum size in polynomial time.

**Solution:** First, we can easily find the size of the minimum vertex cover – call it  $k^*$ . Consider an edge  $u, v$ . Any solution of size  $k^*$  must pick either  $u$  or  $v$ . In other words, either  $G - u$  or  $G - v$  should have a vertex cover of size  $k^* - 1$ . Thus, here is a recursive algorithm  $\mathcal{A}$  which given a graph  $H$  and a number  $k$ , either outputs NO or outputs a vertex cover of size  $k$ . The algorithm  $\mathcal{A}(H, k)$ , where  $k = 1$ , simply checks if there is a single vertex in  $H$  which is a vertex cover. If so, it just returns this vertex, otherwise outputs NO. For  $k > 1$ : let  $(u, v)$  be an edge in  $H$ . Then we recursively run  $\mathcal{A}(H - u, k - 1)$  and  $\mathcal{A}(H - v, k - 1)$ . If the answer is NO in both cases, we return NO. Otherwise say  $\mathcal{A}(H - u, k - 1)$  returns a set  $S$  of size  $k - 1$  which is a vertex cover of  $H - u$ . Then,  $\mathcal{A}(H, k)$  returns  $S \cup \{u\}$  (the other case for  $H - v$  is similar). Finally, we run  $\mathcal{A}$  on  $G, k^*$ .

2. The directed Hamiltonian Cycle Problem is as follows: given a directed graph  $G$ , is there a cycle which contains all the vertices? Suppose you have a polynomial time algorithm for this problem. Show that you can also find such a cycle (if it exists) in polynomial time.

**Solution:** Suppose  $G$  is Hamiltonian. Let  $C$  be any Hamiltonian cycle in  $G$ . Then, if we remove any edge not in  $C$ , the resulting graph will still be Hamiltonian. Thus, we get the following algorithm (let  $\mathcal{A}$  denote the algorithm which given a graph, decides whether it is Hamiltonian or not): first run  $\mathcal{A}$  on  $G$  to check if  $G$  is Hamiltonian or not. Assume  $G$  is Hamiltonian. While  $G$  has more than  $n$  edges, find an edge  $e$  in  $G$  such that  $\mathcal{A}(G - e)$  returns true. As we argued above, there must exist such an edge – so we can try each edge in  $G$  and see if  $\mathcal{A}(G - e)$  is true or not. Let  $e$  be such an edge. Then, we remove  $e$ , and repeat this process. Finally, when  $G$  has only  $n$  edges, these must form a Hamiltonian cycle.

3. The undirected Hamiltonian Cycle Problem can be defined similarly as above. The undirected Hamiltonian Path problem is as follows: given an undirected graph  $G$ , is there a path which contains all the vertices? Show that the undirected Hamiltonian path is polynomial time reducible to the undirected Hamiltonian Cycle problem.

**Solution:** Let  $\mathcal{I}$  be an input to the Hamiltonian path problem. Note that  $\mathcal{I}$  consists of an undirected graph  $G$ . We need to produce a graph  $G'$  such that  $G$  has a Hamiltonian path if and only if  $G'$  has a Hamiltonian cycle. We proceed as follows: add a new vertex  $v$  to the graph  $G$  and add edges between  $v$  and every vertex in  $G$  – call this graph  $G'$ . Now if  $P$  is a Hamiltonian path in  $G$  starting at vertex  $s$  and ending at  $t$ , then

$v, s, P, t, v$  is a Hamiltonian cycle in  $G'$ . Conversely, if  $C$  is a Hamiltonian cycle in  $G'$ , then removing the vertex  $v$  from  $C$  gives a Hamiltonian path in  $G$ .

4. Show that the undirected Hamiltonian cycle problem is reducible to the directed Hamiltonian cycle problem. Show that the directed Hamiltonian cycle problem is reducible to the undirected Hamiltonian cycle problem.

**Solution:** We first reduce the undirected Hamiltonian cycle problem to the directed Hamiltonian cycle problem. Let  $G$  be an undirected graph, which is an input to the undirected Hamiltonian cycle problem. We need to produce a directed graph  $G'$  (in polynomial time) such that  $G$  has a Hamiltonian cycle if and only if  $G'$  has a (directed) Hamiltonian cycle. We construct  $G'$  by replacing each edge in  $G$  by two directed edges (going in opposite direction). It is easy to check that this reduction has the desired property.

The reverse reduction is more tricky. Let  $G = (V, E)$  be a directed graph, and from this we have to produce a graph  $G' = (V', E')$  (in polynomial time) such that  $G$  has a Hamiltonian cycle if and only if  $G'$  has a Hamiltonian cycle. For every vertex  $v \in G$ ,  $G'$  has three vertices -  $v', v'', v'''$  with edges  $(v', v''), (v'', v''')$ . For every directed edge  $(u, v)$  in  $G$ , we have the edge  $(u''', v')$  in  $G'$ . This completes the description of  $G'$ . Now suppose  $G$  has a Hamiltonian cycle:  $v_1, v_2, \dots, v_n$ . Then  $v'_1, v''_1, v'''_1, v'_2, v''_2, v'''_2, \dots$  is a Hamiltonian cycle in  $G'$ . Now, suppose  $G'$  has a Hamiltonian cycle. Since each of the vertices  $v''_i$  has degree 2, they must be preceded by  $v'_i$  and succeeded by  $v'''_i$  in this cycle (or the other way round). Therefore, if the vertices  $v''_i$  appear in the cycle  $C$  in the order  $v''_1, v''_2, \dots, v''_n$ , then it must be the case that the cycle looks like  $v'_1, v''_1, v'''_1, v'_2, v''_2, v'''_2, v'_3, \dots$ , and so  $v_1, v_2, \dots, v_n$ , or  $v_n, v_{n-1}, \dots, v_1$  is a directed cycle in  $G$ .